


## New England Biolabs Certificate of Analysis

**Product Name:** DNase I Reaction Buffer  
**Catalog #:** B0303S  
**Concentration:** 10X Concentrate  
**Lot #:** 0011711  
**Assay Date:** 11/2017  
**Expiration Date:** 11/2020  
**Storage Temp:** -20°C  
**Composition (1X):** 10 mM Tris-HCl, 2.5 mM MgCl<sub>2</sub>, 0.5 mM CaCl<sub>2</sub>, (pH 7.6 @ 25°C)  
**Specification Version:** PS-B0303S v1.0  
**Effective Date:** 16 May 2018

Assay Name/Specification (minimum release criteria)	Lot #0011711
<b>Endonuclease Activity (Nicking, Buffer)</b> - A 50 µl reaction in 1X DNase I Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Functional Testing (DNaseI Reaction Buffer)</b> - A 50 µl reaction in 1X DNase I Reaction Buffer containing 1 µg pBR322 DNA and 1:100 units DNaseI (RNase Free) incubated for 10 minutes at 37°C results in complete digestion of the substrate DNA as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> - A 50 µl reaction in 1X DNase I Reaction Buffer containing 1 µg of HaeIII digested PhiX174 RF I DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>pH (buffers/solutions)</b> - The pH of 10X DNase I Reaction Buffer is between pH 7.5 and 7.7 at 25°C.	<b>Pass</b>
<b>RNase Activity (Buffer)</b> - A 10 µl reaction in 1X DNase I Reaction Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by polyacrylamide gel electrophoresis.	<b>Pass</b>



Authorized by  
Derek Robinson  
16 May 2018



Inspected by  
John Greci  
17 Nov 2017

