

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Thermostable Inorganic Pyrophosphatase

Catalog Number: M0296L
Concentration: 2,000 U/ml

Unit Definition: One unit is the amount of enzyme that will generate 1 µmol of

phosphate per minute from inorganic pyrophosphate under standard reaction conditions (a 10 minute reaction at 75°C in 50 mM Tricine [pH 8.5], 1 mM MgCl2, 0.32 mM PPi, reaction volume of 0.5 ml).

Packaging Lot Number: 10213750
Expiration Date: 10/2025
Storage Temperature: -20°C

Storage Conditions: 20 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

(pH 8.0 @ 25°C)

Specification Version: PS-M0296S/L v1.0

Thermostable Inorganic Pyrophosphatase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0296LVIAL	Thermostable Inorganic Pyrophosphatase	10205127	Pass	

Assay Name/Specification	Lot # 10213750
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 2 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 100 units of Thermostable Inorganic Pyrophosphatase incubated for 4 hours at 75°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 1 containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Thermostable Inorganic Pyrophosphatase incubated for 4 hours at 75°C releases <0.1% of the total radioactivity.	Pass
Phosphatase Activity (pNPP) A 1 ml reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 10 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units of Thermostable Inorganic Pyrophosphatase incubated for 30 minutes at 75°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass



M0296L / Lot: 10213750

Page 1 of 2

Assay Name/Specification	Lot # 10213750
dNTPase Activity	Pass
A 500 µl reaction in CircumVent™ Sequencing Buffer in the presence of 200 µM each	
dNTPs and a minimum of 100 units Thermostable Inorganic Pyrophosphatase incubated	
for 1 hour at 75°C results in <0.01 μmole of inorganic phosphate from dNTPs as	
determined by the AAM assay.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Dongxian Yue Production Scientist 13 Oct 2023 Michael Tonello

Packaging Quality Control Inspector

13 Oct 2023

