

New England Biolabs Certificate of Analysis

Product Name: T3 DNA Ligase
Catalog Number: M0317S
Concentration: 3,000,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to give 50% ligation of 100 ng of Lambda-HindIII fragments in 1 minute at 25°C.
Packaging Lot Number: 10094830
Expiration Date: 01/2023
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0317S/L v2.0


T3 DNA Ligase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0317SVIAL	T3 DNA Ligase	10094829	Pass
B0535AVIAL	StickTogether™ DNA Ligase Buffer	10077408	Pass

Assay Name/Specification	Lot # 10094830
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of T3 DNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 15000 units of T3 DNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 1 containing 1 µg of CIP-treated Lambda-HindIII DNA and a minimum of 3000 units of T3 DNA Ligase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity, Digested DNA)	Pass

Assay Name/Specification	Lot # 10094830
<p>A 20 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 2 µg of Lambda DNA-HindIII Digest and a minimum of 3,000 units of T3 DNA Ligase incubated for 16 hours at 37°C results in >95% ligation of the DNA fragments as determined by agarose gel electrophoresis. Of these ligated fragments, >95% can be recut with HindIII.</p>	
<p>Functional Testing (Adaptor Ligation) A 20 µl reaction in 1X StickTogether™ DNA Ligase Buffer containing 40 µM of phosphorylated linker and 3,000 units of T3 DNA Ligase incubated for 16 hours at 16°C results in no detectable unligated adaptor as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) T3 DNA Ligase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>Protein Concentration (A280) The concentration of T3 DNA Ligase is 1 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 62,130 and molecular weight of 39,351 daltons for T3 DNA Ligase (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423).</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 15000 units of T3 DNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
22 Feb 2021



Michael Tonello
Packaging Quality Control Inspector
22 Feb 2021