240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: LongAmp® Taq DNA Polymerase

Catalog #: M0323S/L
Concentration: 2,500 units/ml

Unit Definition: One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes

at 75°C.

 Lot #:
 0161606

 Assay Date:
 06/2016

 Expiration Date:
 06/2018

 Storage Temp:
 -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween® 20, 0.5 % IGEPAL® CA-630, 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0323S/L v1.0

Effective Date: 13 Jun 2016

Assay Name/Specification (minimum release criteria)	Lot #0161606
Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in NEBuffer 2 containing 1 μg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2.5 units of LongAmp® <i>Taq</i> DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (30 kb Human Genomic DNA) - A 25 μl reaction in LongAmp® <i>Taq</i> Reaction Buffer in the presence of 300 μM dNTPs and 0.4 μM primers containing 500 ng Human Genomic DNA with 2.5 units of LongAmp® <i>Taq</i> DNA Polymerase for 28 cycles of PCR amplification results in the expected 30 kb product.	Pass
PCR Amplification (30 kb Lambda DNA) - A 25 μl reaction in LongAmp® <i>Taq</i> Reaction Buffer in the presence of 300 μM dNTPs and 0.4 μM primers containing 1 ng Lambda DNA with 2.5 units of LongAmp® <i>Taq</i> DNA Polymerase for 28 cycles of PCR amplification results in the expected 30 kb product.	Pass
<b>qPCR DNA Contamination (</b> <i>E. coli</i> <b>Genomic)</b> - A minimum of 2.5 units of LongAmp® $Taq$ DNA Polymerase is screened for the presence of <i>E. coli</i> genomic DNA using SYBR® Green qPCR with primers specific for the <i>E. coli</i> 16S rRNA locus. Results are quantified using a standard curve generated from purified <i>E. coli</i> genomic DNA. The measured level of <i>E. coli</i> genomic DNA contamination is $\leq 1$ <i>E. coli</i> genome.	Pass







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0161606
RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-	
stranded RNA and a minimum of 1 µl of LongAmp® Taq DNA Polymerase is incubated at 37°C. After incubation	Pass
for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent	F 455
detection.	

M.W. Southworth

Authorized by Maurice Southworth 13 Jun 2016







Inspected by
Cathy Rezac
21 Jun 2016