

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: T4 RNA Ligase 2, truncated KQ

Catalog Number: M0373S
Concentration: 200,000 U/ml

Unit Definition: 200 units is defined as the amount of enzyme required to give 80%

ligation of a 31-mer RNA to the pre-adenylated end of a 17-mer DNA

in a total reaction volume of 20 µl in 1 hour at 25°C.

Packaging Lot Number: 10127450
Expiration Date: 09/2023
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 10 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0373S/L v2.0

T4 RNA Ligase 2, truncated KQ Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0373SVIAL	T4 RNA Ligase 2, truncated KQ	10122070	Pass	
B1004SVIAL	PEG 8000	10115452	Pass	
B0216SVIAL	T4 RNA Ligase Reaction Buffer	10122381	Pass	

Assay Name/Specification	Lot # 10127450
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in T4 RNA Ligase Reaction Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 200 units of T4 RNA Ligase 2, truncated KQ incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 μl reaction in T4 RNA Ligase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 200 units of T4 RNA Ligase 2, truncated KQ incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 200 units of T4 RNA Ligase 2, truncated KQ is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass



M0373S / Lot: 10127450 Page 1 of 2

Assay Name/Specification	Lot # 10127450
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 200 units of T4 RNA Ligase 2, truncated KQ incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE) T4 RNA Ligase 2, truncated KQ is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist

02 Dec 2021

Michael Tonello

Packaging Quality Control Inspector

02 Dec 2021



M0373S / Lot: 10127450

Page 2 of 2