

New England Biolabs Certificate of Analysis

Product Name: Rapid™ PNGase F
Catalog #: P0710S
Concentration: 1 reaction/μl
Unit Definition: N/A
Lot #: 0021802
Assay Date: 02/2018
Expiration Date: 02/2019
Storage Temp: 4°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)
Specification Version: PS-P0710S v2.0
Effective Date: 17 Jan 2018

| Assay Name/Specification (minimum release criteria) | Lot #0021802 |
|---|--------------|
| Glycosidase Activity (Endo F1, F2, H) - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (Endo F2, F3) - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β-Mannosidase) - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β-Xylosidase) - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β1-3 Galactosidase) - A 10 μl reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β-Galactosidase substrate (Galβ1-3GlcNAcβ1-4Galβ1-4Glc-AMC) and 1 μl of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |



New England Biolabs Certificate of Analysis

| Assay Name/Specification (minimum release criteria) | Lot #0021802 |
|---|--------------|
| Glycosidase Activity (β1-4 Galactosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β -Galactosidase substrate (Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc -AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β-N-Acetylgalactosaminidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β -N-Acetylgalactosaminidase substrate (GalNAc β 1-4Gal β 1-4Glc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (β-N-Acetylglucosaminidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled β -N-Acetylglucosaminidase substrate (GlcNAc β 1-4GlcNAc β 1-4GlcNAc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α-Glucosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Glucosidase substrate (Glc α 1-6Glc α 1-4Glc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α-Neuraminidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Neuraminidase substrate (Neu5Ac α 2-3Gal β 1-3GlcNAc β 1-3Gal β 1-4Glc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α1-2 Fucosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-2Gal β 1-4Glc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α1-3 Fucosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Fucosidase substrate (Fuc α 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4Glc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α1-3 Galactosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-3Gal β 1-4GlcNAc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α1-3 Mannosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-3Man β 1-4GlcNAc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |

New England Biolabs Certificate of Analysis

| Assay Name/Specification (minimum release criteria) | Lot #0021802 |
|--|--------------|
| Glycosidase Activity (α1-6 Galactosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Galactosidase substrate (Gal α 1-6Gal α 1-6Glc α 1-2Fru-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α1-6 Mannosidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -Mannosidase substrate (Man α 1-6Man α 1-6(Man α 1-3)Man-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Glycosidase Activity (α-N-Acetylgalactosaminidase) - A 10 μ l reaction in Rapid PNGase F Buffer containing 1 nM of fluorescently-labeled α -N-Acetylgalactosaminidase substrate (GalNAc α 1-3(Fuc α 1-2)Gal β 1-4Glc-AMC) and 1 μ l of Rapid PNGase F incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography. | Pass |
| Protease Activity (SDS-PAGE) - A 20 μ l reaction in 1X Rapid PNGase F Buffer containing 24 μ g of a standard mixture of proteins and a minimum of 5 μ l of Rapid PNGase F incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection. | Pass |
| Protein Purity Assay (SDS-PAGE) - Rapid PNGase F is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |



Authorized by
Derek Robinson
17 Jan 2018



Inspected by
Alicia Bielik
06 Apr 2018

