

New England Biolabs Certificate of Analysis

Product Name: Proteinase K, Molecular Biology Grade
Catalog Number: P8107S
Concentration: 800 U/ml
Unit Definition: One unit will digest urea-denatured hemoglobin at 37°C (pH 7.5) per minute to produce equal absorbance as 1.0 µmol L-tyrosine using Folin & Ciocalteu's phenol reagent.
Packaging Lot Number: 10061037
Expiration Date: 11/2021
Storage Temperature: -20°C
Storage Conditions: 20 mM Tris-HCl, 1 mM CaCl₂, 50% Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-P8107S v1.0

Proteinase K, Molecular Biology Grade Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
P8107SVIAL	Proteinase K, Molecular Biology Grade	10058455	Pass


Assay Name/Specification	Lot # 10061037
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>qPCR DNA Contamination (Eukaryotic Genomic) A minimum of 1.6 units of Proteinase K, Molecular Biology Grade is screened for the presence of eukaryotic genomic DNA using SYBR® Green qPCR with universal primers for the 18S rRNA locus. Results are quantified using a standard curve generated from purified E. album genomic DNA. The measured level of eukaryotic genomic DNA contamination is ≤ 2.5 pg DNA/µl.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo)</p>	Pass

Assay Name/Specification	Lot # 10061037
<p>A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 4 units of Proteinase K, Molecular Biology Grade incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 8 units of Proteinase K, Molecular Biology Grade incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 RF I DNA and a minimum of 0.8 units of Proteinase K, Molecular Biology Grade incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Beth Paschal
Production Scientist
01 Nov 2019



Jay Minichiello
Packaging Quality Control Inspector
10 Jan 2020