

New England Biolabs Certificate of Analysis

Product Name: HgaI
Catalog #: R0154S/L
Concentration: 2,000 units/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 DNA in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot #: 0291301
Assay Date: 01/2013
Expiration Date: 01/2015
Storage Temp: -20 °C
Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0154S/L v1.0
Effective Date: 20 Jun 2013

Assay Name/Specification (minimum release criteria)	Lot #0291301
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 2 units of HgaI incubated for 4 hours at 37°C releases <0.2% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) - After a 2-fold over-digestion of PhiX174 DNA with HgaI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with HgaI.	Pass
Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 1.1 containing 1 µg of PhiX174 DNA and a minimum of 2 Units of HgaI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) - HgaI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by
Derek Robinson
20 Jun 2013



Inspected by
Leo Formenoy
20 Jun 2013

