

New England Biolabs Certificate of Analysis

Product Name: Goat Anti-Rabbit IgG Magnetic Beads
Catalog #: S1432S
Concentration: 20 mg/ml
Lot #: 0101711
Assay Date: 11/2017
Expiration Date: 11/2020
Storage Temp: 4°C
Storage Conditions: 0.02 % NaN₃, 0.05 % Tween®20, 0.1 % BSA, 1 X PBS, (pH 7.4 @ 25°C)
Specification Version: PS-S1432S v1.0
Effective Date: 15 Jun 2018

| Assay Name/Specification (minimum release criteria) | Lot #0101711 |
|---|--------------|
| Binding Capacity (Magnetic Beads) - Goat Anti-Rabbit IgG Magnetic Beads (1 mg) were equilibrated and incubated with 200 µl of >0.5 mg/ml Purified Rabbit IgG for 1 hour at 25°C, then washed and the Rabbit IgG eluted. Binding capacity was determined to be >5 µg of Rabbit IgG per mg of beads. | Pass |
| Functional Binding Assay (Qualitative) - Goat Anti-Rabbit IgG Magnetic Beads (1 mg) were equilibrated and incubated with 500 µl of Serum containing Rabbit IgG for 1 hour at 25°C, then washed, eluted and evaluated by Tris-Glycine gel to confirm low non-specific binding of extract proteins and high isolation of target. | Pass |
| Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in Goat Anti-Rabbit IgG Magnetic Bead Storage Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| RNase Activity (Buffer) - A 10 µl reaction in Goat Anti-Rabbit IgG Magnetic Bead Storage Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection. | Pass |



Authorized by
Derek Robinson
15 Jun 2018



Inspected by
Michael Sproviero
03 Nov 2017

