

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:	Goat Anti-Rat IgG Magnetic Beads
Catalog #:	<i>S1433S</i>
Concentration:	20 mg/ml
Lot #:	0051804
Assay Date:	04/2018
Expiration Date:	04/2021
Storage Temp:	4°C
Storage Conditions:	0.02 % NaN3, 0.05 % Tween® 20, 0.1 % BSA, 1 X PBS, (pH 7.4 @ 25°C)
Specification Version:	PS-S1433S v1.0
Effective Date:	15 Jun 2018

Assay Name/Specification (minimum release criteria)	Lot #0051804 Pass
Binding Capacity (Magnetic Beads) - Goat Anti-Rat IgG Magnetic Beads (1 mg) were equilibrated and incubated with 200 μ l of >0.5 mg/ml Purified Rat IgG for 1 hour at 25°C, then washed and the Rat IgG eluted. Binding capacity was determined to be >5 μ g of Rat IgG per mg of beads.	
Functional Binding Assay (Qualitative) - Goat Anti-Rat IgG Magnetic Beads (1 mg) were equilibrated and incubated with 500 µl of Serum containing Rat IgG for 1 hour at 25°C, then washed, eluted and evaluated by Tris -Glycine gel to confirm low non-specific binding of extract proteins and high isolation of target.	
Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in Goat Anti-Rat IgG Magnetic Bead Storage Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
RNase Activity (Buffer) - A 10 μ l reaction in Goat Anti-Rat IgG Magnetic Bead Storage Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by fluorescent detection.	

un otun

Authorized by Derek Robinson 15 Jun 2018



Inspected by Michael Sproviero 01 Apr 2018