

New England Biolabs Product Specification

<i>Product Name:</i>	<i>Tth Endonuclease IV</i>
<i>Catalog #:</i>	<i>M0294S</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to cleave 1 pmol of a 60-mer oligonucleotide duplex containing a single AP site in a total reaction volume of 10 µl in 1 hour at 65°C.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 % Triton®X-100, 50 % Glycerol, (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0294S v2.0</i>
<i>Effective Date:</i>	<i>12 Aug 2019</i>

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of Tth Endonuclease IV incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer 1 containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of Tth Endonuclease IV incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of Tth Endonuclease IV incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.



Date 12 Aug 2019

Derek Robinson
Director of Quality Control

